

There might be superluminal particles in nature *

Guang-jiong Ni [†]

Department of Physics, Fudan University, Shanghai, 200433, P.R.China

Department of Physics, Portland State University, Portland, OR 97207 USA.

Abstract

Based on experimental discovery that the mass-square of neutrino is negative, a quantum equation for superluminal neutrino is proposed in comparison with Dirac equation and Dirac equation with imaginary mass. A virtual particle may also be viewed as superluminal one. Both the basic symmetry of space-time inversion and the maximum violation of space-inversion symmetry are emphasized.

Key words: ordinary neutrinos; neutrino mass and mixing; non-standard-model neutrinos

1 INTRODUCTION

Nothing can travel faster than light. Is this statement as true now as it ever was? In 2000 there were two experiments showing superluminal propagation of microwave ^[1] or laser pulse ^[2]. Yet physicists believe that the law of physics have remained intact ^[3]. However, the experimental discovery of negative mass square of neutrino in recent years ^[4],

$$E^2 = c^2 p^2 + m^2 c^4, \quad (1)$$

$$m^2(\nu_e) = -2.5 \pm 3.3 eV^2, \quad (2)$$

though far from accurate, does strongly hint that neutrino might be a particle moving faster than light. Actually, rewriting (1) as

$$E^2 = c^2 p^2 - m_s^2 c^4 \quad (3)$$

and using the quantum relations $E = \hbar\omega$ and $p = \hbar k$, one easily derives the kinematic relation for superluminal particle (also named as tachyon in literature) as follows (m_s is called "proper mass"):

$$p = \tilde{m}u = \frac{m_s u}{\sqrt{\frac{u^2}{c^2} - 1}}, \quad E = \tilde{m}c^2 = \frac{m_s c^2}{\sqrt{\frac{u^2}{c^2} - 1}}. \quad (4)$$

Here the velocity of particle u is identified with the group velocity $u_g = \frac{d\omega}{dk}$ of wave. To derive (3), a quantum equation for neutrino was proposed in Ref. [5,6] (see also [7]). In this paper, we will compare three kinds of equation—Dirac equation and two "superluminal" equations and discuss relevant problems.

*The paper has been published in Journal of Shaanxi Normal University (Natural Science Edition), Vol.29, No.3

[†]E-mail: gj_ni@Yahoo.com

2 Dirac equation versus superluminal equation

The Dirac equation for fermion with rest mass m_0 reads

$$i\hbar \frac{\partial}{\partial t} \psi = i\hbar \vec{\alpha} \cdot \nabla \psi + \beta m_0 c^2 \psi \quad (5)$$

where ψ is a four-component spinor wave function and α_i and β are 4×4 matrixes. In Dirac representation, they are expressed as

$$\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}, \quad \alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \quad (6)$$

where φ and χ are two-component spinors, σ_i are Pauli matrixes. Now we perform a unitary transformation as

$$\psi \longrightarrow U\psi = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \psi = \psi' = \begin{pmatrix} \xi \\ \eta \end{pmatrix}. \quad (7)$$

Let the matrix U act on Eq.(5) from the left, due to noncommutativity between U and α_i (or β), we find Dirac equation in "Weyl representation" with

$$\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad \alpha_i^{(W)} = \begin{pmatrix} \sigma_i & 0 \\ 0 & -\sigma_i \end{pmatrix}, \quad \beta^{(W)} = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \quad (8)$$

Note that, however, the representation transformation leads to important change in physical interpretation. While φ and χ in (6) represent the "hidden particle and antiparticle fields" in a particle state^[8], ξ and η in (8) characterize the "hidden left-handed and right-handed rotating fields" in a particle with 100% left-handed helicity explicitly.

Now let's perform a unitary transformation on Dirac equation (5) in Weyl representation:

$$\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix} \longrightarrow U_s \psi = \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix} \psi = \psi_s = \begin{pmatrix} \xi_s \\ \eta_s \end{pmatrix}. \quad (9)$$

After setting

$$m_s = -im_0, \quad (10)$$

we obtain an equation with real proper mass m_s :

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \xi_s &= i\hbar \vec{\sigma} \cdot \nabla \xi_s - m_s c^2 \eta_s, \\ i\hbar \frac{\partial}{\partial t} \eta_s &= -i\hbar \vec{\sigma} \cdot \nabla \eta_s + m_s c^2 \xi_s. \end{aligned} \quad (11)$$

A substitution of plane-wave solution $\xi_s \sim \eta_s \sim \exp[i(px - Et)/\hbar]$ leads to Eq. (3) immediately. Why so simple transformation like (9) can change Dirac equation into a "superluminal" equation? It is because the U_s in (9) is a nonhermitian matrix. While a unitary but hermitian matrix like that in (7) merely amounts to a representation transformation, a unitary but nonhermitian matrix is capable of changing the physical essence of equation. Eq.(11) can be expressed in Dirac representation as

$$i\hbar \frac{\partial}{\partial t} \psi_s = i\hbar \vec{\alpha} \cdot \nabla \psi_s + \beta_s m_s c^2 \psi_s \quad (12)$$

$$\psi_s = \begin{pmatrix} \varphi_s \\ \chi_s \end{pmatrix} = U \begin{pmatrix} \xi_s \\ \eta_s \end{pmatrix}, \quad \beta_s = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}. \quad (13)$$

Note that the matrix β_s and its counterpart in Eq. (11) are antihermitian.

On the other hand, if we directly set $m_0 = im_s$ in Eq. (5) to get the "Dirac equation with imaginary mass" as

$$i\hbar \frac{\partial}{\partial t} \psi^{(i)} = ic\hbar \vec{\alpha} \cdot \nabla \psi^{(i)} + \beta_s^{(i)} m_s c^2 \psi^{(i)}, \quad (14)$$

then we have

$$\psi^{(i)} = \begin{pmatrix} \varphi^{(i)} \\ \chi^{(i)} \end{pmatrix}, \quad \beta_s^{(i)} = \begin{pmatrix} iI & 0 \\ 0 & -iI \end{pmatrix}. \quad (15)$$

The fact that equation (14) is wrong can easily be seen by its plane-wave solution when $p \rightarrow 0$, yielding a decoupling solution:

$$\begin{aligned} \varphi^{(i)} &\sim e^{-i(im_s t)} \sim e^{m_s t} \\ \chi^{(i)} &\sim e^{-i(-im_s t)} \sim e^{-m_s t} \end{aligned} \quad (16)$$

which show the violation of unitarity. By contrast, such kind of meaningless solution is definitely excluded in Eq. (12). The sharp difference is stemming from the following fact. Three kinds of Eqs. (5), (12) and (14), all respect the basic symmetry: under the space-time inversion ($\vec{x} \rightarrow -\vec{x}$, $t \rightarrow -t$) [8]

$$\varphi(-\vec{x}, -t) \rightarrow \chi(\vec{x}, t), \quad \chi(-\vec{x}, -t) \rightarrow \varphi(\vec{x}, t), \quad (17)$$

the theory remains invariant (while a concrete solution of particle transforms into that of antiparticle). However, we should consider a smaller symmetry—the space-inversion ($\vec{x} \rightarrow -\vec{x}$, $t \rightarrow t$):

$$\xi(-\vec{x}, t) \rightarrow \eta(\vec{x}, t), \quad \eta(-\vec{x}, t) \rightarrow \xi(\vec{x}, t). \quad (18)$$

Then we see that Eqs. (5) and (14) remain invariant whereas Eq. (12) (or (11)) fails to do so. In other words, our new superluminal equation reflects the maximum parity violation, a property exactly explaining the permanent helicity of neutrino—while ν_L and $\bar{\nu}_R$ are physically realized, ν_R and $\bar{\nu}_L$ are forbidden strictly—as verified by experiments [9,10].

3 Virtual particle as a superluminal particle

It is no surprise to see solution like (16) when Eq. (15) contains an imaginary mass im_s explicitly. Actually, in quantum mechanics, we often endow an imaginary part of mass m to wave function for describing an unstable particle:

$$m \rightarrow m - i\frac{\Gamma}{2}, \quad \Gamma = \hbar/\tau, \quad |\psi|^2 \sim e^{-t/\tau} \quad (19)$$

with τ being the decay lifetime. Of course, the unitarity of wave function is destroyed.

Hence, it is hopeless to set $m_0 = im_s$ directly in Dirac equation for describing a stable superluminal particle. However, we often discuss the "virtual particle" in covariant perturbation theory in the sense that its momentum p and energy E can vary independently and so are not subjected to the constraint $E^2 - p^2 = m_0^2$. Both $E^2 > p^2$ and $E^2 < p^2$ cases can occur. We might as well say that a virtual particle could be superluminal but it is unstable too. A stable superluminal particle can only be realized as ν_L or $\bar{\nu}_R$ ensured by the violation of parity symmetry.

4 Maximum parity violation displays the beauty of nature

Usually, a symmetry, i.e., an invariance under some transformation yields an affirmative guarantee for a theory and its violation seems to spoil the validity of the theory. Now we see that for a meaningful discrete symmetry like space-inversion, what happens in nature is either keeping intact (for subluminal particle) or being violated to maximum (for superluminal neutrino). In the latter case the violation must reach its maximum because the constraint of larger symmetry—the space-time invariance—must be held at the same time. [In the first line of Eq. (11), the coefficient of m_s must be fixed as (-1) to ensure the invariance of transformation (17) together with the violation of transformation (18).] We might as well look at the maximum violation of parity as some "antisymmetry" (rather than "asymmetry") which also provides affirmative guarantee for the validity of superluminal theory for neutrino. In fact, the normal consequence of violation of hermitian property being the violation of unitarity (as shown by (16)) is now recast into a strange realization—of four would be unstable solutions for a same momentum, two of them (ν_R and $\bar{\nu}_L$) are eliminated whereas other two (ν_L and $\bar{\nu}_R$) are stabilized.

5 Summary and discussion

We can have three kinds of Dirac-type equation:

$$i\hbar\frac{\partial}{\partial t}\psi = i\hbar\vec{\alpha}\cdot\nabla\psi + \beta mc^2\psi \quad (20)$$

as listed in the following table.

Finally, two remarks are in order:

(a) According to our present understanding, no boson but fermion can be superluminal as long as the parity symmetry is violated to maximum. Most likely, one kind of known particles, the neutrino, is just a superluminal particle, a tachyon with spin 1/2.

(b) The physical implication of nonhermitian transformation U_s amounts to an extra phase difference $\pi/2$ between ξ_s and η_s in comparison with that in subluminal case. Hence we see once again that it is the phase which plays a dominant role in quantum mechanics.

(c) The origin (or mechanism) of mass generation has been remained in physics as a mystery for hundred years. It is time to say something now. According to detailed analysis in Ref. [6], the finite and changeable mass of a fermion is not due to one kind of excitation which varies continuously inside but a manifestation of coherent cancellation effect between two fields rotating with opposite helicities implicitly, either one of them can be excited infinitely in essence. It is precisely this fantastic effect explains the amazing result that a tachyon's energy (mass) approaches zero when its velocity u increases to infinity (as shown in Eq. (4)).

Acknowledgements: The author wishes to thank Dr. Tsao Chang for bringing the superluminal problem of neutrino to his attention and relevant discussions.

References

- [1] D. Mugnai, A. Ranfagni and R. Ruggeri, Observation of superluminal behaviors in wave propagation, Phys. Rev. Lett. 84(2000)4830.
- [2] L. J. Wang, A. Kuzmich and A. Dogariu, Gain-assisted superluminal light propagation, Nature, 406, 20 July, (2000)277.
- [3] A. M. Steinberg, Nothing goes faster than light, Physics World, Sep. 2000,21-21.
- [4] "Review of Particle Physics", Euro. Phys. J. C 15 (2000) 350.
- [5] T. Chang and G.-j. Ni, An explanation on negative mass-square of neutrinos, Preprint, hep-ph/0009291.
- [6] G.-j. Ni and T. Chang, Is neutrino a superluminal particle? Preprint, hep-ph/0103051.
- [7] J.Ciborowski and J.Rembielinski, Tritium decay and the Hypothesis of tachyonic neutrinos, Euro. Phys. J., 1999,C8:157.
- [8] G.-j. Ni, H. Guan, W.-m. Zhou and J. Yan, Antiparticle in the light of Einstein-Podolsky-Rosen paradox and Klein paradox, Chin. Phys. Lett. 17(2000) 393.
- [9] T.D. Lee and C.N. Yang, Question of parity conservation in weak interactions, Phys. Rev. 104 (1956) 254.
- [10] C.S. Wu E.Ambler, R.W.Hayward, et al, Experimental test of parity conservation in beta decay, Phys. Rev. 105 (1957) 1413.

property	Dirac equation	Superluminal equation Eq. (11), (12)	Dirac equation with imaginary mass Eq. (14)
m (mass)	m_0 (rest mass)	m_s (proper mass)	m_s
Dirac representation $\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}, \quad \vec{\alpha} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}$	$\beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$ hermitian matrix	$\beta_s = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ antihermitian matrix	$\beta_s^{(i)} = \begin{pmatrix} iI & 0 \\ 0 & -iI \end{pmatrix}$ antihermitian matrix
Weyl representation $\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad \vec{\alpha} = \begin{pmatrix} \vec{\sigma} & 0 \\ 0 & -\vec{\sigma} \end{pmatrix}$	$\beta = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$ hermitian matrix	$\beta_s = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ antihermitian matrix	$\beta_s^{(i)} = \begin{pmatrix} 0 & iI \\ iI & 0 \end{pmatrix}$ antihermitian matrix
hermitian property of theory	yes	no	no
unitarity of theory	yes	yes	no
invariance under space-time inversion (basic symmetry)	yes	yes	yes
invariance under space inversion (parity)	yes	no	yes
physical meaning	subluminal particles (electron, etc.)	describing possibly the (stable) superluminal neutrino ν_L and $\bar{\nu}_R$ (with ν_R and $\bar{\nu}_L$ forbidden)	unstable superluminal particle without physical meaning in reality